

# Proof of Euler's Formula

Adrian D'Costa

We have to prove that:

$$e^{i\theta} = \cos(\theta) + i \sin(\theta)$$

Let  $f(\theta) = e^{i\theta}$  According to Maclaurin series we know that:

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)x^n}{n!} \quad \text{where } f^{(0)}(a) = f(a) \text{ and } 0! = 1$$

Also:

$$i^2 = -1, i^3 = -i, i^4 = 1, i^5 = i, i^6 = -1, i^7 = -i, i^8 = 1$$

$$\begin{aligned} f^{(0)}(0) &= 1 \\ f^{(1)}(0) &= i \\ f^{(2)}(0) &= -1 \\ f^{(3)}(0) &= -i \\ f^{(4)}(0) &= 1 \\ f^{(5)}(0) &= i \\ f^{(6)}(0) &= -1 \\ f^{(7)}(0) &= -i \\ f^{(8)}(0) &= 1 \end{aligned}$$

$$e^{i\theta} = \frac{1}{0!} + \frac{i\theta}{1!} - \frac{\theta^2}{2!} - \frac{i\theta^3}{3!} + \frac{\theta^4}{4!} + \frac{i\theta^5}{5!} - \frac{\theta^6}{6!} - \frac{i\theta^7}{7!} + \frac{\theta^8}{8!} + \dots$$

Now to find Maclaurin series representation of:

$$g(\theta) = \cos(\theta)$$

$$\begin{aligned} g^{(0)}(0) &= \cos(0) = 1 \\ g^{(1)}(0) &= -\sin(0) = 0 \\ g^{(2)}(0) &= -\cos(0) = -1 \\ g^{(3)}(0) &= \sin(0) = 0 \\ g^{(4)}(0) &= \cos(0) = 1 \\ g^{(5)}(0) &= -\sin(0) = 0 \\ g^{(6)}(0) &= -\cos(0) = -1 \\ g^{(7)}(0) &= \sin(0) = 0 \\ g^{(8)}(0) &= \cos(0) = 1 \end{aligned}$$

$$\cos(\theta) = \frac{1}{0!} - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \frac{\theta^6}{6!} + \frac{\theta^8}{8!} + \dots$$

Maclaurin series representation of:

$$h(\theta) = \sin(x)$$

$$\begin{aligned}
h^{(0)}(0) &= \sin(0) = 0 \\
h^{(1)}(0) &= \cos(0) = 1 \\
h^{(2)}(0) &= -\sin(0) = 0 \\
h^{(3)}(0) &= -\cos(0) = -1 \\
h^{(4)}(0) &= \sin(0) = 0
\end{aligned}$$

$$\begin{aligned}
h^{(5)}(0) &= \cos(0) = 1 \\
h^{(6)}(0) &= -\sin(0) = 0 \\
h^{(7)}(0) &= -\cos(0) = -1 \\
h^{(8)}(0) &= \sin(0) = 0
\end{aligned}$$

$$i \sin(\theta) = \frac{i\theta}{1!} - \frac{i\theta^3}{3!} + \frac{i\theta^5}{5!} - \frac{i\theta^7}{7!} + \dots$$

Now:

$$\begin{aligned}
\cos(\theta) + i \sin(\theta) &= \frac{1}{0!} - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \frac{\theta^6}{6!} + \frac{\theta^8}{8!} + \frac{i\theta}{1!} - \frac{i\theta^3}{3!} + \frac{i\theta^5}{5!} - \frac{i\theta^7}{7!} + \dots \\
&= \frac{1}{0!} + \frac{i\theta}{1!} - \frac{\theta^2}{2!} - \frac{i\theta^3}{3!} + \frac{\theta^4}{4!} + \frac{i\theta^5}{5!} - \frac{\theta^6}{6!} - \frac{i\theta^7}{7!} + \frac{\theta^8}{8!} + \dots \\
&= e^{i\theta}
\end{aligned}$$

[Q.E.D.]

## 0.1 Another way to prove Euler's formula

Another way to prove Euler's formula is that:

$$\cos(\theta) + i \sin(\theta) = e^{i\theta}$$

Now rearranging that equation we get:

$$1 = \frac{e^{i\theta}}{\cos(\theta) + i \sin(\theta)}$$

$$0 = \frac{d}{d\theta} \left( \frac{e^{i\theta}}{\cos(\theta) + i \sin(\theta)} \right) \dots \text{[Taking der. of both sides]}$$

$$0 = \frac{ie^{i\theta} \cos(\theta) - e^{i\theta} \sin(\theta) + e^{i\theta} \sin(\theta) - ie^{i\theta} \cos(\theta)}{\cos^2(\theta) + 2i \cos(\theta) \sin(\theta) - \sin^2(\theta)}$$

$$0 = \frac{0}{\cos^2(\theta) + 2i \cos(\theta) \sin(\theta) - \sin^2(\theta)}$$

$$0 = 0$$

$$L.H.S = R.H.S$$

[Q.E.D.]